



EUROPEAN COMMISSION  
JOINT RESEARCH CENTRE

Directorate F - Health, Consumers and Reference Materials (Geel)  
Food and Feed Compliance



JRC F.5/CvH/MGH/AS/Ares

**Evaluation Report on the Analytical Methods submitted  
in connection with the Application for Authorisation of a  
Feed Additive according to Regulation (EC) No 1831/2003**

**Monosodium L-glutamate**  
produced by fermentation with  
*Corynebacterium glutamicum* KCCM80187  
(FAD-2019-0055; CRL/190032)





**Evaluation Report on the Analytical Methods submitted  
in connection with the Application for Authorisation of a  
Feed Additive according to Regulation (EC) No 1831/2003**

Dossier related to: **FAD-2019-0055 - CRL/190032**

Name of Product: ***Monosodium L-glutamate  
produced by fermentation with  
Corynebacterium glutamicum  
KCCM80187***

Active Agent (s): **Monosodium L-glutamate**

Rapporteur Laboratory: **European Union Reference Laboratory for  
Feed Additives (EURL-FA)  
JRC Geel, Belgium**

Report prepared by: **María José González de la Huebra**

Report checked by: **Zigmas Ezerskis**  
Date: **15/02/2021**

Report approved by: **Christoph von Holst**  
Date: **16/02/2021**

## EXECUTIVE SUMMARY

In the current application an authorisation is sought under Article 4(1) for *monosodium L-glutamate (MSG)* produced by fermentation with *Corynebacterium glutamicum* KCCM80187, under the category/functional group 2(b) 'sensory additives'/flavouring compounds', according to Annex I of Regulation (EC) No 1831/2003. The authorisation is sought for all animal species.

According to the Applicant *MSG* has a minimum purity (mass fraction) of 99 %. The *feed additive* is intended to be added directly into *feedingstuffs* (or through *premixtures*) and *water* for drinking. The Applicant proposed a recommended maximum content of *MSG* in *feedingstuffs* of 25 mg/kg.

For the quantification of *MSG* in the *feed additive* the Applicant submitted an in-house validated analytical method based on reversed phase high performance liquid chromatography coupled with ultraviolet detection (HPLC-UV). While in the frame of the validation study satisfactory performance characteristics were derived, the Applicant did not present a verification study or any additional test performed by a second independent laboratory applying the above mentioned method.

For the quantification of *MSG* in *premixtures* and *feedingstuffs* the Applicant submitted the ring-trial validated European Union method (Commission Regulation (EC) No 152/2009) based on ion-exchange chromatography coupled to photometric detection (IEC-VIS). This method, designed only for the analysis of amino acids in *premixtures* and *feedingstuffs*, does not distinguish between the amino acids and their salts, or between different salts of the same amino acids, and it cannot differentiate between enantiomers.

The method was further ring-trial validated resulting in the EN ISO 13903:2005 method. The following performance characteristics were reported for the quantification of glutamic acid: a relative standard deviation for *repeatability* (RSD<sub>r</sub>) ranging from 0.9 to 2.7 % and a relative standard deviation for *reproducibility* (RSD<sub>R</sub>) ranging from 6.2 to 9.1 %. However, while a lowest limit of quantification (LOQ) of 30 mg/kg has been reported for the analysis of certain amino acids, a specific LOQ for glutamic acid has not been indicated. Therefore the method does not ensure the accurate determination of *MSG* when added into feed at the proposed recommended maximum content (i.e. 25 mg/kg *feedingstuffs*).

Hence, the EURL recommends for official control the European Union method based on IEC-VIS for the quantification of *MSG* in *premixtures* only. The Applicant did not provide any experimental data to determine *MSG* in *water*. Nevertheless, as concluded in previous EURL reports on amino acids, the EURL recommends for official control the procedure based on IEC-VIS and described in the ring-trial validated European Union method (or in equivalent

ring-trial validated methods e.g. VDLUFA Method 4.11.6.) to quantify *MSG* in in the *feed additive* and *water*.

The methods described above do not distinguish between the amino acids and their salts, or between different salts of the same amino acids, and they cannot differentiate between enantiomers. As a consequence, *MSG* is detected as glutamic acid.

In addition, the EURL recommends the "Monosodium L-glutamate monograph" of the Food Chemical Codex (FCC) for the identification of the *feed additive*.

Further testing or validation of the methods to be performed through the consortium of National Reference Laboratories as specified by Article 10 (Commission Regulation (EC) No 378/2005, as last amended by Regulation (EU) 2015/1761) is not considered necessary.

## KEYWORDS

*Monosodium L-glutamate (MSG)* produced by fermentation with *Corynebacterium glutamicum* KCCM80187, glutamic acid, sensory additives, flavouring compounds, all animal species.

## 1. BACKGROUND

In the current application an authorisation is sought under Article 4(1) (authorisation of a new feed additive) for *monosodium L-glutamate (MSG)* produced by fermentation with *Corynebacterium glutamicum* KCCM80187, under the category/functional group 2(b) 'sensory additives'/flavouring compounds', according to Annex I of Regulation (EC) No 1831/2003. The authorisation is sought for all animal species [1-2]. Synthetically produced *MSG* is already authorised as sensory *feed additive* [3-4].

According to the Applicant, *MSG* (monosodium salt of glutamic acid) is an off-white crystalline powder with a minimum purity (mass fraction) of 99 % [5]. The *feed additive* is produced by fermentation with a non-toxic, non-pathogenic species of *Corynebacterium glutamicum* KCCM80187 [6]. The production strain is deposited in the "Korean Culture Centre of Microorganisms" (KCCM) with the accession number 80187 [7].

The *feed additive* is intended to be added directly into *feedingstuffs* or through *premixtures* and *water* for drinking [8]. Furthermore, the Applicant proposed a recommended maximum content of *MSG* in *feedingstuffs* of 25 mg/kg [1,8].

Note: The EURL previously evaluated the analytical methods for the determination of *MSG* as sensory *feed additive* in the frame of previous dossiers [9].

## 2. TERMS OF REFERENCE

In accordance with Article 5 of Regulation (EC) No 378/2005, as last amended by Regulation (EU) 2015/1761, on detailed rules for the implementation of Regulation (EC) No 1831/2003 of the European Parliament and of the Council as regards the duties and the tasks of the European Union Reference Laboratory concerning applications for authorisations of feed additives, the EURL is requested to submit a full evaluation report to the European Food Safety Authority for each application or group of applications. For this particular dossier, the methods of analysis submitted in connection with *monosodium L-glutamate (MSG)* produced by fermentation with *Corynebacterium glutamicum* KCCM80187 and their suitability to be used for official controls in the frame of the authorisation were evaluated.

## 3. EVALUATION

***Description of the analytical methods for the determination of the active substance in the feed additive, premixtures, feedingstuffs and when appropriate water (section 2.6.1 of the dossier - Annex II of Commission Regulation (EC) No 429/2008)***

For the quantification of *MSG* in the *feed additive* the Applicant submitted an in-house validated analytical method based on reversed phase high performance liquid chromatography (HPLC) and ultraviolet detection (UV) [10,11].

The *feed additive* is diluted in distilled water and an aliquot is directly injected into the HPLC system equipped with a C18 column. The analyte is eluted via a potassium phosphate buffer. *MSG* is detected with a photodiode array detector (PDA) at 210 nm and quantified via a calibration curve. In the frame of the validation study, the Applicant reported relative standard deviations for *repeatability* ( $RSD_r$ ) and *intermediate precision* ( $RSD_{ip}$ ) below 3 %, and a *recovery rate* ( $R_{rec}$ ) ranging from 99 to 105 % [11]. While in the frame of the validation study satisfactory performance characteristics were derived, the Applicant did not present a verification study or any additional test performed by a second independent laboratory applying the above mentioned method.

For the quantification of the *MSG* content in *premixtures* and *feedingstuffs* the Applicant submitted the ring-trial validated European Union method [10,12]. This method applies for the determination of free (synthetic and natural) and total (peptide-bound and free) amino acids (including glutamic acid), using an amino acid analyser or HPLC equipment with an ion exchange column (IEC). The method is intended for *premixtures* and *feedingstuffs*, it does not distinguish between the amino acids and their salts, or between different salts of the same amino acids and it cannot differentiate between enantiomers. As a consequence, *MSG* is detected as glutamic acid.

The *free* amino acids are extracted with diluted hydrochloric acid. Co-extracted nitrogenous macromolecules are precipitated with sulfosalicylic acid and removed by filtration. The solution is filtered and adjusted to pH 2.2. The amino acids are separated by IEC and *free* glutamic acid is determined after post-column derivatisation with ninhydrin by spectrophotometric detection at 570 nm (visible-VIS).

The procedure chosen for the determination of the *total* amino acids depends on the amino acids under investigation. Glutamic acid can be determined in either oxidised or non-oxidised samples. Oxidation is performed at 0 °C with a performic acid/phenol mixture. The excess of the oxidation reagent is decomposed with sodium disulfite. The oxidised or non-oxidised sample is hydrolysed with hydrochloric acid (6 mol/l) containing 1 g phenol/l for 23 hours. The hydrolysate is adjusted to pH 2.2. The amino acids are separated by IEC and total glutamic acid is determined by post-column derivatisation with ninhydrin and photometric detection at 570 nm.

As a general rule, in order to avoid an overestimation of added *MSG* content, the procedure based on quantification of *free* amino acids content in *premixtures* and *feedingstuffs* has to be followed. When the procedure for the quantification of *total* amino acids is applied on *premixtures* and *feedingstuffs* containing proteins, an overestimation of *MSG* may occur.

The European Union method was further ring-trial validated by twenty-three laboratories for the determination of total glutamic acid in *feed* and resulted in the equivalent standard method EN ISO 13903:2005 [13]. The reported performance characteristics are listed in Table 1.

While limits of quantification (LOQ) ranging from 30 to 350 mg/kg have been reported for the analysis of free amino acids, a specific LOQ for glutamic acid has not been indicated [13]. However, according to the EURL's former evaluations, this method does not ensure the accurate discrimination between *MSG* as flavouring added at low concentration to *feedingstuffs* (i.e. 25 mg/kg) and endogenous *MSG*/glutamic acid possibly present at higher concentration levels in the *feed* matrix [9]. Therefore the EURL is unable to recommend a method for the official control to determine *MSG* added into *feedingstuffs*.

**Table 1:** Method performance characteristics obtained in the frame of EN ISO 13903:2005 [13] for the determination of total *glutamic acid* in *feed*.

Ring-Trial	Matrix	Glutamic acid content g/kg	RSD <sub>r</sub> %	RSD <sub>R</sub> %
[13]	Poultry meal	79.7	2.7	9.1
	Broiler finisher feed	32.5	1.6	7.0
	Broiler starter feed	40.4	1.8	8.4
	Corn	15.1	2.4	6.2
	Fishmeal	73.7	0.9	4.7

RSD<sub>r</sub>, RSD<sub>R</sub> - relative standard deviation for *repeatability* and *reproducibility*, respectively.

Based on the performance characteristics presented, the EURL recommends for official control the ring-trial validated European Union method based on IEC-VIS to quantify *MSG* in *premixtures*.

The Applicant did not provide any experimental data to determine *MSG* in *water* [10]. Nevertheless, based on NRL's experience, as concluded in previous EURL's amino acids reports [14] even if the determination of *MSG* in the *feed additive* and *water* is not explicitly stated in the scope of the European Union method (or similar ones e.g. VDLUFA Method 4.11.6.), the IEC-VIS procedure described above is considered fit-for-purpose for the determination of *MSG* in these matrices.

Therefore, the EURL recommends for official control the procedure based on IEC-VIS and described in the ring-trial validated European Union method to quantify *MSG* in the *feed additive* and *water*.

***Methods of analysis for the determination of the residues of the additive in food (section 2.6.2 of the dossier - Annex II of Commission Regulation (EC) No 429/2008)***

The evaluation of corresponding methods of analysis is not relevant for the present application.

***Identification/Characterisation of the feed additive (section 2.6.3 of the dossier - Annex II of Commission Regulation (EC) No 429/2008)***

For the identification of the *feed additive* the EURL recommends the "Monosodium L-glutamate monograph" of the Food Chemical Codex (FCC) where different tests (including one for the specific optical rotation) are described [15].

Further testing or validation of the methods to be performed through the consortium of National Reference Laboratories as specified by Article 10 (Commission Regulation (EC) No 378/2005, as last amended by Regulation (EU) 2015/1761) is not considered necessary.

## **4. CONCLUSIONS AND RECOMMENDATIONS**

In the frame of this authorisation the EURL recommends for official control (i) the "Monosodium L-glutamate monograph" of the Food Chemical Codex (FCC) for the identification of *MSG* in the *feed additive*; (ii) the European Union method based on IEC-VIS (or equivalent method) for the quantification of *MSG* in the *feed additive* and *water*; and (iii) the European Union method based on IEC-VIS for the quantification of *MSG* in *premixtures*.



***Recommended text for the register entry (analytical method)***

For the identification of *monosodium L-glutamate* in the *feed additive*:

- Food Chemical Codex "Monosodium L-glutamate monograph"

For the quantification of *monosodium L-glutamate* in the *feed additive* and *water*:

- ion-exchange chromatography coupled with post-column derivatisation and photometric detection (IEC-VIS)

For the quantification of *monosodium L-glutamate* in *premixtures*:

- ion-exchange chromatography coupled with post-column derivatisation and photometric detection (IEC-VIS), Commission Regulation (EC) No 152/2009 (Annex III, F)

**5. DOCUMENTATION AND SAMPLES PROVIDED TO EURL**

In accordance with the requirements of Regulation (EC) No 1831/2003, reference samples of *monosodium L-glutamate* produced by fermentation with *Corynebacterium glutamicum* KCCM80187 have been sent to the European Union Reference Laboratory for Feed Additives. The dossier has been made available to the EURL by EFSA.

**6. REFERENCES**

- [1] \* Application, Reference SANTE/E5: Forw. Appl. 1831/0001-2020
- [2] \* Annex I – submission number 1566215097276-2438
- [3] Commission Implementing Regulation (EU) 2018/249 of 15 February 2018 concerning the authorisation of taurine, beta-alanine, L-alanine, L-arginine, L-aspartic acid, L-histidine, D,L-isoleucine, L-leucine, L-phenylalanine, L-proline, D,L-serine, L-tyrosine, L-methionine, L-valine, L-cysteine, glycine, monosodium glutamate and L-glutamic acid as feed additives for all animal species and L-cysteine hydrochloride monohydrate for all species except cats and dogs, O.J. L 53/134, 23.2.2018
- [4] Commission Implementing Regulation (EU) 2018/1567 of 18 October 2018 correcting Implementing Regulation (EU) 2018/249 concerning the authorisation of taurine, beta-alanine, L-alanine, L-arginine, L-aspartic acid, L-histidine, D,L-isoleucine, L-leucine, L-phenylalanine, L-proline, D,L-serine, L-tyrosine, L-methionine, L-valine, L-cysteine, glycine, monosodium glutamate and L-glutamic acid as feed additives for all animal species and L-cysteine hydrochloride monohydrate for all species except cats and dogs O.J. L 262/32, 19.10.2018
- [5] \*Technical dossier, Section II: II.2.1.1. Chemical substances
- [6] \*Technical dossier, Section II: II.2.1.2. Micro-organisms
- [7] \*Technical dossier, Section II: II.1.1. Name of the additive
- [8] \*Technical dossier, Section II: II.5.1. Proposed mode of use in animal nutrition

- [9] EURL evaluation Reports:  
<https://ec.europa.eu/jrc/sites/jrcsh/files/FinRep-FAD-2010-0107.pdf>  
[https://ec.europa.eu/jrc/sites/jrcsh/files/finrep\\_fad-2018-0090-msglutamate.pdf](https://ec.europa.eu/jrc/sites/jrcsh/files/finrep_fad-2018-0090-msglutamate.pdf)
- [10] \*Technical dossier, Section II: II.6.1. Methods of analysis for the active substance
- [11] \*Technical dossier, Section II: Annex\_II\_6\_01 CJ MSG Method validation report.pdf
- [12] Commission Regulation (EC) No 152/2009 of 27 January 2009 laying down the methods of sampling and analysis for the official control of feed, O.J. L 54, 26.02.2009
- [13] EN ISO 13903:2005- Animal feeding stuffs – Determination of amino acids content
- [14] EURL evaluation Report:  
[https://ec.europa.eu/jrc/sites/jrcsh/files/finrep\\_fad-2018-0045\\_l-arginine.pdf](https://ec.europa.eu/jrc/sites/jrcsh/files/finrep_fad-2018-0045_l-arginine.pdf)
- [15] Food Chemical Codex monograph "monosodium L-glutamate", FCC 7 (2010), p.698
- \*Refers to Dossier no: FAD-2019-0055

## 7. RAPPORTEUR LABORATORY & NATIONAL REFERENCE LABORATORIES

The Rapporteur Laboratory for this evaluation is the European Union Reference Laboratory for Feed Additives, JRC, Geel, Belgium. This report is in accordance with the opinion of the consortium of National Reference Laboratories as referred to in Article 6(2) of Commission Regulation (EC) No 378/2005, as last amended by Regulation (EU) 2015/1761.

## 8. ACKNOWLEDGEMENTS

The following National Reference Laboratories contributed to this report:

- Centro di referenza nazionale per la sorveglianza ed il controllo degli alimenti per gli animali (CReAA), Torino (IT)
- Państwowy Instytut Weterynaryjny, Pulawy (PL)
- Österreichische Agentur für Gesundheit und Ernährungssicherheit (AGES), Wien (AT)
- Thüringer Landesanstalt für Landwirtschaft (TLL). Abteilung Untersuchungswesen. Jena (DE)
- Laboratoire de Rennes (SCL L35), Service Commun des Laboratoires DGCCRF et DGDDI, Rennes (FR)
- Wageningen Food Safety Research<sup>1</sup> (WFSR) (NL)
- Ústřední kontrolní a zkušební ústav zemědělský (ÚKZÚZ), Praha (CZ)

---

<sup>1</sup> Name and address according to according COMMISSION IMPLEMENTING REGULATION (EU) 2015/1761: RIKILT Wageningen UR, Wageningen.