

Evolution of Copernicus Land Services based on Sentinel data

2019 IACS Workshop

CLMS-related H2020 projects session

Wednesday, 10th April 2019

08:30 – 18:00 CET

Valladolid (Spain), Consejería de Agricultura y Ganadería JCYL

ECoLass



ECoLaSS: "Evolution of Copernicus Land Services based on Sentinel data"

Eva Sevillano Marco, Katharina Schwab, Markus Probeck (GAF)














H2020 ECoLass – Partners involved

ECoLass: "Evolution of Copernicus Land Services based on Sentinel data"

Key Objective = **improve** existing & develop **novel** products/services for future operational pan-European & Global Copernicus Land Components 2020+:










services in the land mapping domain



H2020 ECoLass – Time scale

ECoLaSS concept within the time lapse of Copernicus phases development and the Copernicus Land Monitoring Service frame





Dates & Timing


Dec 2015 Horizon 2020 WP.

Call: "Evolution of Copernicus Land Monitoring Services"

Jan 2017: Project Start

→ Runtime: Jan 2017 - Dec 2019 (3 years)















H2020 ECoLass - Objectives & Setup

OBJECTIVES

- Innovative methods: improve existing & develop novel products and services
- Prototypes: next-generation operational Copernicus Land Services demonstrated in large prototype sites of biogeographical diversity + food security
- Operationalization framework: benchmarking of prototypes for operational implementation into Copernicus Land services +2020



High data volume processing chains

Service Requirements



✓ User Needs

✓ EO and Other Data

✓Infrastructure & Architecture

Methods



✓ Sentinel Optical+SAR Integration

√ Time Series Analysis, Classification & Change

✓ Incremental HRL Updates

Prototypes



✓ Next-generation Services/ Products for Continental & Global Component

✓ Demonstrated on Large Sites (Europe & Africa)

Operationalization



✓ Stakeholder Consultation & Endorsement /

√Operational Maturity

✓ Integration into Copernicus Land 2020+



Method developments for dense time series









H2020 ECoLass – Objectives & Setup

Implementation

Task 2 User Requirements

Stakeholders surveys

- Thematic topics & Service evolution
- Earth Observation & In-situ data
- Architecture & Infrastructure



Task 5 Operationalization Framework

Prototypes benchmarking for implementation into Copernicus Land Monitoring Services 2020+ Innovation & Technical excellence





Task 3 Innovative Methods


Time series Sentinel optical+SAR

- Pre-processing & fusion
- Classification
- Change detection
- Incremental updates



Task 4 Prototypes

Demonstration in large sites of biogeographical diversity



Africa














H2020 ECoLass – Challenges & Issues addressed

Copernicus Land Monitoring Service



Multiresolution Multisensor Multitemporal

Challenges & Issues

 HR local-continental-global scales o data gaps (e.g., cloud cover) oproducts consistency

Research needs

LC/LU and change mapping **Multisensor** data (radar + optical) Automated change monitoring based on Sentine S Improved **grasslands** identification methods Crop area and crop status monitoring Methods to provide yearly incremental updates in HRL **High volume data** processing lines



EO data **Sentinel TS 2014-present** spatial information on various spatial scales biophysical variables LC/LU status +change layers



Evolving user requirements and state of the art R&D methodologies driving Copernicus operational services

> R&D results support informed discussion of Copernicus service portfolio









H2020 ECoLass – Outcomes

Requirements

Service evolution

- o Not much thought/interest about technical issues, specifications
- High interest in Copernicus satellite data (mostly Sentinel-2)
- Higher interest in local component/higher resolution

EO & Reference data

Input EO data


- Shortcomings VHR data: unequal European DTM, geometric accuracy
- o S1 image frame position inconsistencies due to other radar interference
- S2 geometric and cloud masks inconsistencies

In-situ data

- CORDA is huge improvement
- Heterogeneity for thematic classification (e.g.≈ IACS European coverage)

IT Infrastructure

- Network/data transfer, storage, processing, scalability capabilities
- o Processing techniques will favour solutions that move software to data
- o DIAS still not yet operationally mature











H2020 ECoLass – Outcomes

Requirements

Service Evolution Requirements in view of potential +2020 implementation

- o state of the art
- product specifications
- identifying challenges
- o shortcomings and lessons learned

CLMS global component

Existing: MR-HR Sentinel layers

New: phenology layers

CLMS local component

Existing: Natura 2000, Urban Atlas

New: Coastal Zone Service

CLMS Pan-European component

Existing:

HRLs qualitative improvements & more frequent updates, CLC+ spatial resolution and thematic content

New: Agriculture Service, phenology layer, generic products

Higher resolution Frequent updates Thematic content **Timeliness**
















H2020 ECoLass – Outcomes Methods & tests

Methods development & tests


Larger prototype sites (≈100 000 km²) demonstration of candidates for Copernicus Land Service Evolution roll-out on a larger scale

Test sites with specific **in-situ** data ○ **IACS-LUCAS-HRL**-others

Europe



S2 tile ≈100x100 km



Africa



o JECAM (Joint Experiment for Crop Assessment and Monitoring) test sites in Belgium, France, Mali and South Africa ensure large crop and field size diversity, including grasslands and natural vegetation.














Background: High Resolution Layers (HRLs) 2015 → **ECoLaSS Prototypes**












Forest


Status layers 10 m

- o Dominant Leaf Type (broadleaved/coniferous) 2017, 2018
- Tree Cover Density 2018

Change products 20 m

Forest Cover Change (FCC) 2015-2017 & 2017-2018





- -North, Central, and South-East demosites
- -Sentinel-1 + Sentinel-2 data
- -Reference data
- o HRLs 2015
- o VHR 2015 & 2017/2018










Imperviousness

© Google. Image © 2017 DigitalGlobe

© European Union, Copernicus Land Monitoring Service 2015, European Environment Agency (EEA).


Status layers 10 m

Improved imperviousness degree 2017, 2018



PA = 92,4% (+/-1,6%) UA = 85,1% (+/-1,7%) Omission = 7,6%Commission = 14,9%

2017 Belgium



Change products 20 m

o Incremental imperviousness change 2015-2017 & 2017-2018









Grasslands

S2 Grassland masks tests 2018

S1+S2

Grassland masks tests 2018


S1+S2

Status layers 10 m

- o Grassland mask 2017, 2018
- Use intensity 2018

Change products 20 m

o Grassland change 2015-2017 & 2017-2018



Class	HRL GRA 2015	ECoLaSS grassland
		prototype 2017
Grassland	10,365 km² / 83.9 %	10,943 km² / 83.2 %
Non-Grassland	53,957 km² / 16.1 %	54,083 km² / 16.8 %
Clouds/Cloud-	$2 \mathrm{km^2}$ / 0.0%	$0\mathrm{km^2}$ / 0.0%
Shadows		
Overall	64,324 km ²	65,023 km ²

© European Union, Copernicus Land Monitoring Service 2015, European Environment Agency (EEA).













Legend

Maize

Peas

Chicory

Winter Cereals

Summer Wheat

Hemp

Green Beans

Onions

Fruit Crops

Other Crops
Grape Vines

Not classified Grassland Winter Wheat

Winter Barley
Sugar Beet
Spelt
Rapeseed
Potato
Flax

Summer Oat
Other Vegetables
Other Fodder

Summer Barley Summer Cereals

Agriculture

Mali, Belgium/France, Central demosites


- -Sentinel-1 + Sentinel-2 TS
- -Reference data: IACS, HRLs 2015, LUCAS,
- Africa In-situ field data collected

© Google. Image © 2018 DigitalGlobe

Crop mask 2018

Crop Mask 2016, 2017, 2018 Crop type masks 2017, 2018

Crop types 2017







Status layers 10 m












New Land Cover

New land cover layer at 10m, with 7 classes that is being modified to include CLC+ recommendations



- OA = 94%
- PA = 82-99%
- UA = 89-97%
- Kappa = 0.8
- F-Score = 0.37-0.48
- Data 1 year coverage of S2 data only (yearly updates)
 Production Improved automation for faster production
 Definition Thematic harmonization between pan-European and global layers



Crop, grasslands and forest masks combination

In ECoLaSS the prototypes include across products consistency by testing combination between HRLs










Indicators

© Google. Image © 2017 DigitalGlobe



Experimental prototypes related to Indicators and Variables


- Crop growth condition
- Crop Emergence Date Map
- Generic Land Cover Metrics


Maximum phenology layer (Max month NDVI)

Multi-annual trends & potential change

Start of season (March, April, May)

End of season (August, September, October, Winter)





Peak of season (May, June, July, August)

Length of season (6, 7, 8, 9 months)

Crop growth condition S2 2017 Belgium test











H2020 ECoLass – Evaluation candidate prototypes

Benchmarking

Methods & prototypes

- Sentinel-1/2/3 integration

- Sentinel/TS Pre-processing

- TS classification

- TS change detection

- Incremental updates

Benchmarking Approach



High-volume data processing chains: automated, flexible & customizable



Prototypes:

- Indicators and variables
- HRL incremental updates
- Grassland characterisation
- Crop status/monitoring
- New Products

Operational roll-out candidates

	Crop growth condition	New product New product New product		
Time series indicators	Crop Emergence Date Map			
Time series indicators	Generic Land Cover Metrics			
	Multi-annual trends & potential change	New product		
Imperviousness incremental update	Improved IMD status layer at 10m	Improved product		
Imperviousness incremental appate	Incremental IMD change	Improved product		
Forest incremental update	Improved DLT status layer at 10m	Improved product		
Porest incremental update	Incremental tree cover (mask) change	Improved product		
Improved Grassland	Improved Grassland status layer at 10m	Improved product		
No. Action the constant	New crop mask status layer at 10m	New product		
New Agriculture product	New crop type status layer at 10m	New product		
New Products	CLC evolution	New product		

Assessment matrix

- long term evolution
- o portfolio complementarities
- o answering identified needs
- o political support
- o state of the art
- maturity/timing

- EO & in-situ availability
- processing capacity
- automation level
- o roll-out potential
- cost/benefit
- documentation











Evaluation candidate prototypes

Assessment matrix

s	ervice/product candidate	long-term evolution	portfolio comple- mentarity	answering identified needs	political support	respecting core vs. DS	State of the art/Innovatio	Maturity/ Timing	adequate EO availability	adequate in- situ availability	processing capacity (platform, SW)	Automation level	pratically proven Roll- out potential	Cost/ benefit (forecast)	Documen- tation	Overall score
ir	nproved IMD status layer at 10m		o	+	++	++	+	++	++	+	++	++	+	+	+/++	59,50
ir	cremental IMD change	++	+	++	++	++	++	+	+	o	++	+	+	+	+/++	61,50
ir	nproved DLT status layer at 10m		0	+	#	++	+	++	++	+	++	+	+	+	+/++	58,50
ir	cremental forest cover change	++	+	++	+	++	o/+	+	+	o	++	+	+	+	+/++	60,00
	nproved Grassland status layer at Om	0	0	++	+	++	+	++	++	+	++	+	+	+	+/++	60,50
	ew crop mask status layer at 10m	++	+	++		++	++	o	**	-	++	+	+	+	+/++	58,50
N N	ew crop type status layer at 10m	++	+	++		++	++	o	++		++	+	o	+	+/++	56,50
c	LC evolution	+	+	++	+	++	++	o	++	o	o	o	-	+	+	54,00
	rop growth condition	++	+	+	0	o	++	+	++		++	++	+	+	+	56,00
	rop Emergence Date Map	+	+	0	0	o	++	0	+		++	+	+	+	+/++	52,50
G	eneric Land Cover Metrics	++	+	+	+	+/++	+	-	+	o	++	o	-	+	0	51,50
	Julti-annual trends & potential nange	++	+	0	0	+/++	+		++	0	++	o/+	-/o	+	0	51,50





++	very satisfactory/relevant/applicable					
+	satisfactory/relevant/applicable					
0	neutral					
-	not satisfactory/relevant/applicable					
	not at all satisfactory/relevant/applicable					











Evaluation candidate prototypes

CONCLUSIONS

✓ The prototypes **improve** the **Copernicus HRLs** and develop new products, fulfilling identified requirements



✓ Benchmarking analysis evaluate prototypes against manifold criteria (e.g. long-term evolution, maturity/timing, automation level, cost/benefit (forecast)).

	Copernicus HRL	ECoLaSS HRL
Spatial Resolution	20m	10m
Temporal Resolution	3 years	1 year
Methodology	manual & automated processing	mostly automated
Themes	Imperviousness	Imperviousness
	Forests	Forests
	Grassland	Grassland
	Water & Wetness	
	Landcover	Landcover
		Temporal Indicators
		Agriculture

Score	Verdict			
61.50				
60.50 most promisin				
60.00				
58.50				
59.50				
58.50	high potential			
56.50				
56.00				
54.00				
52.50				
51.50	experimental			
51.50				
	61.50 60.50 60.00 58.50 59.50 58.50 56.50 56.00 54.00 52.50 51.50			

- √HRL Status layers 10 m/ Change layers 20 m
- ✓ **Updates** propel services **continuous** production: yearly update feasible through automated TS exploitation
- ✓Incorporating **phenological** and **statistical parameters** to accurate masks, continuous scales are derived for a tailored management
- ✓ Challenges: **timeliness** production for operational services at pan-European and global scales in a proper spatial resolution & products **consistency** (status, change, updates)









Follow the ECoLaSS team !!

www.ecolass.eu Follow us:




GAFAG L. Moser, M. Probeck, K. Schwab, G. Ramminger, E.Sevillano & Team

C. Sannier, S. Villerot, B. Desclée, A. Masse & Development Team

H. Gallaun, M. Schardt, P. Miletich, J. Deutscher, K. Granica

P. Defourny, E. Khalil, J. Wolter, I. Moreau, X. Blaes

B. Leutner, A. Metz-Marconcini, I. Klein, A. Hirner, T. Esch





The research leading to these results has received funding from the European Union's Horizon 2020 Research and Innovation Programme, under Grant Agreement n° 730008.





Horizon2020











